

Product Datasheet

NM23-H1 Antibody (13C2-R)

NBP3-33132

Unit Size: 100 ul

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NBP3-33132

Updated 8/7/2025 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NBP3-33132



NBP3-33132

NM23-H1 Antibody (13C2-R)

Product Information	
Unit Size	100 ul
Concentration	1 mg/ml
Storage	Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.
Clonality	Monoclonal
Clone	13C2-R
Preservative	0.05% Sodium Azide
Isotype	IgG1
Purity	Protein A purified
Buffer	PBS (pH7.4), 0.1% BSA and 40% Glycerol
Target Molecular Weight	17 kDa
Product Description	
Host	Mouse
Gene ID	4830
Gene Symbol	NME1
Species	Human, Mouse, Rat
Immunogen	Synthetic peptide within Human NM23-H1 aa 1-50 / 152. (Uniprot: P15531)
Product Application Details	
Applications	Western Blot, Immunohistochemistry-Paraffin, Immunocytochemistry/Immunofluorescence, Immunohistochemistry
Recommended Dilutions	Western Blot 1:2000, Immunohistochemistry, Immunocytochemistry/Immunofluorescence 1:100, Immunohistochemistry-Paraffin 1:2000



Images

Western Blot: NM23-H1 Antibody (13C2-R) [NBP3-33132] - Western blot analysis of NM23-H1 on different lysates with Mouse anti-NM23-H1 antibody (NBP3-33132) at 1/2,000 dilution.

Lane 1: HEK-293 cell lysate (20 ug/Lane)
 Lane 2: MCF7 cell lysate (20 ug/Lane)
 Lane 3: Jurkat cell lysate (20 ug/Lane)
 Lane 4: Raji cell lysate (20 ug/Lane)
 Lane 5: HeLa cell lysate (20 ug/Lane)
 Lane 6: K-562 cell lysate (20 ug/Lane)
 Lane 7: A549 cell lysate (20 ug/Lane)
 Lane 8: A431 cell lysate (20 ug/Lane)
 Lane 9: HepG2 cell lysate (20 ug/Lane)
 Lane 10: NIH/3T3 cell lysate (20 ug/Lane)
 Lane 11: C6 cell lysate (20 ug/Lane)
 Lane 12: Mouse kidney tissue lysate (40 ug/Lane)
 Lane 13: Mouse liver tissue lysate (40 ug/Lane)
 Lane 14: Rat brain tissue lysate (40 ug/Lane)

Predicted band size: 17 kDa
 Observed band size: 17/20 kDa



Exposure time: 1 minute 59 seconds;

4-20% SDS-PAGE gel.

Proteins were transferred to a PVDF membrane and blocked with 5% NFDM/TBST for 1 hour at room temperature. The primary antibody (NBP3-33132) at 1/2,000 dilution was used in 5% NFDM/TBST at 4C overnight. Goat Anti-Mouse IgG - HRP Secondary Antibody at 1/50,000 dilution was used for 1 hour at room temperature.

Immunohistochemistry: NM23-H1 Antibody (13C2-R) [NBP3-33132] - Immunohistochemical analysis of paraffin-embedded mouse kidney tissue with Mouse anti-NM23-H1 antibody (NBP3-33132) at 1/2,000 dilution.


The section was pre-treated using heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 20 minutes. The tissues were blocked in 1% BSA for 20 minutes at room temperature, washed with ddH₂O and PBS, and then probed with the primary antibody (NBP3-33132) at 1/2,000 dilution for 1 hour at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.



Immunocytochemistry/ Immunofluorescence: NM23-H1 Antibody (13C2-R) [NBP3-33132] - Immunocytochemistry analysis of HeLa cells labeling NM23-H1 with Mouse anti-NM23-H1 antibody (NBP3-33132) at 1/100 dilution.

Cells were fixed in 4% paraformaldehyde for 20 minutes at room temperature, permeabilized with 0.1% Triton X-100 in PBS for 5 minutes at room temperature, then blocked with 1% BSA in 10% negative goat serum for 1 hour at room temperature. Cells were then incubated with Mouse anti-NM23-H1 antibody (NBP3-33132) at 1/100 dilution in 1% BSA in PBST overnight at 4 °C. Goat Anti-Mouse IgG H&L (iFluor™ 488) was used as the secondary antibody at 1/1,000 dilution. PBS instead of the primary antibody was used as the secondary antibody only control. Nuclear DNA was labelled in blue with DAPI.

beta Tubulin (red) was stained at 1/100 dilution overnight at +4 °C. Goat Anti-Rabbit IgG H&L (iFluor™ 594) were used as the secondary antibody at 1/1,000 dilution.





Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NBP3-33132

HAF007	Goat anti-Mouse IgG Secondary Antibody [HRP]
NB720-B	Rabbit anti-Mouse IgG (H+L) Secondary Antibody [Biotin]
NBP1-97005-0.5mg	Mouse IgG1 Isotype Control (MG1)
NBP2-13662PEP	NM23-H1 Recombinant Protein Antigen

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP3-33132

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

