# **Product Datasheet**

## Integrin alpha 5 beta 1 Antibody (SG/19) [Alexa Fluor® 405] - Chimeric NBP3-43579AF405

Unit Size: 0.1 ml

Store at 4C in the dark.

www.novusbio.com

technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP3-43579AF405

Updated 3/24/2025 v.20.1

Earn rewards for product reviews and publications.



Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP3-43579AF405

#### NBP3-43579AF405

Integrin alpha 5 beta 1 Antibody (SG/19) [Alexa Fluor® 405] - Chimeric

Product InformationUnit Size0.1 mlConcentrationPlease see the vial label for concentration. If unlisted please contact technical services.StorageStore at 4C in the dark.ClonalityMonoclonalCloneSG/19Preservative0.05% Sodium AzideIsotypeIgGConjugateAlexa Fluor 405PurityProtein A purifiedBuffer50mM Sodium BorateProduct Description3678Gene ID3678SpeciesHumanSpeciesThis antibody binds human integrin alpha 5 bet 1 near the N-terminus. Epitope mapping of SG/19 revealed that Thr82 in the 1 subunit, located at the outer face of the boundary between the 1-like and proteinases and proteinases and thereby stimulates angiogenesis. Integrin alpha5 beta 1 is also known as fibronectin receptor and binds matrix macromolecules and proteinases and thereby stimulates angiogenesis. Integrin alpha5 beta 1 is also known as fibronectin receptor and binds matrix macromolecules and proteinases and thereby stimulates angiogenesis. Integrin alpha5 beta 1 is also known as fibronectin receptor and binds matrix macromolecules and proteinases and thereby stimulates angiogenesis. Integrin alpha5 beta 1 is also known as fibronectin receptor and binds matrix macromolecules and proteinases and thereby stimulates angiogenesis. Integrin alpha5 beta 1 is also known as fibronectin receptor and binds matrix macromolecules and proteinases and thereby stimulates angiogenesis. Integrin alpha5 beta 1 is also known as fibronectin receptor and binds matrix macromolecules and proteinases and thereby stimulates angiogenesis. Integrin alpha5 beta 1 is also known as fibronectin receptor and binds matrix macromolecules and proteinases and thereby stimulates angiog		
ConcentrationPlease see the vial label for concentration. If unlisted please contact technical services.StorageStore at 4C in the dark.ClonalityMonoclonalCloneSG/19Preservative0.05% Sodium AzideIsotypeIgGConjugateAlexa Fluor 405PurityProtein A purifiedBuffer50mM Sodium BorateProduct DescriptionRabbitGene ID3678Gene SymbolITGA5SpeciesHumanSpecificity/SensitivityThis antibody binds human integrin alpha 5 beta 1 near the N-terminus. Epitope mapping SG/19 revealed that Thr82 in the 1 subunit, located at the outer face of the boundary between the I-like and hybrid domains, was the key binding determinant for this antibody. Integrin alpha5 beta 1 is also known as fibronectin receptor and binds matrix macromolecules and proteinases and thereby stimulates angiogenesis. Integrin alpha5 beta 1 is also known as fibronectin family, is a heterodimer composed by two subunits, alpha5 (ITGA5/CD49e) and beta1 (ITGB1/CD29), and both are necessary for complete biological functions. The interaction of VLA-5 with fibronectin plays an important role in regulating inflammatory cytokine production by human articular chondrocytes (From the Cell Migration Gateway ITGA5 ITGB1ImmunogenThe original antibody was generated by immunizing BALB/c mice with Nalm-6	Product Information	
services.StorageStore at 4C in the dark.ClonalityMonoclonalCloneSG/19Preservative0.05% Sodium AzideIsotypeIgGConjugateAlexa Fluor 405PurityProtein A purifiedBuffer50mM Sodium BorateProduct DescriptionRabbitGene ID3678Gene SymbolITGA5SpeciesHumanSpecificity/SensitivityThis antibody binds human integrin alpha 5 beta 1 near the N-terminus. Epitope mapping of SG/19 revealed that Thr82 in the 1 subunit, located at the outer face of the boundary between the I-like and hybrid domains, was the key binding determinant for this antibody. Integrin alpha5beta 1, is a member of the integrin family, is a heterodimer composed by two subunits, alpha5 (ITGA5/CD49e) and beta1 (ITGB1/CD29), and both are necessary for complete biological functions. The interaction of VLA-5 with fibronectin plays an important role in regulating inflammatory cytokine production by human articular chondrocytes (From the Cell Migration Gateway ITGA5 ITGB1ImmunogenThe original antibody was generated by immunizing BALB/c mice with Nalm-6	Unit Size	0.1 ml
ClonalityMonoclonalCloneSG/19Preservative0.05% Sodium AzideIsotypeIgGConjugateAlexa Fluor 405PurityProtein A purifiedBuffer50mM Sodium BorateProduct DescriptionHostRabbitGene ID3678Gene SymbolITGA5SpeciesHumanSpecificity/SensitivityThis antibody binds human integrin alpha 5 beta 1 near the N-terminus. Epitope mapping of SG/19 revealed that Thr82 in the 1 subunit, located at the outer face of the boundary between the 1-like and hybrid domains, was the key binding determinant for this antibody. Integrin alpha5 beta 1 is also known as fibronectin receptor and binds matrix macromolecules and proteinases and thereby stimulates angiogenesis. Integrin alpha5 (ITGA5/CD49e) and beta1 (ITGB1/CD29), and both are necessary for complete biological functions. The interaction of VLA-5 with fibronectin plays an important role in regulating inflammatory cytokine production by human articular chondrocytes (From the Cell Migration Gateway ITGA5 ITGB1ImmunogenThe original antibody was generated by immunizing BALB/c mice with Nalm-6	Concentration	•
CloneSG/19Preservative0.05% Sodium AzideIsotypeIgGConjugateAlexa Fluor 405PurityProtein A purifiedBuffer50mM Sodium BorateProduct DescriptionHostRabbitGene ID3678Gene SymbolITGA5SpeciesHumanSpecificity/SensitivityThis antibody binds human integrin alpha 5 beta 1 near the N-terminus. Epitope mapping of SG/19 revealed that Thr82 in the 1 subunit, located at the outer face of the boundary between the I-like and hybrid domains, was the key binding determinant for this antibody. Integrin alpha5 beta 1 is also known as fibronectin receptor and binds matrix macromolecules and proteinases and thereby stimulates angiogenesis. Integrin alpha5beta1 is a member of the integrin family, is a heterodimer composed by two subunits, alpha5b(ITGA5/CD49e) and beta1 (ITGB1/CD29), and both are necessary for complete biological functions. The interaction of VLA-5 with fibronectin plays an important role in regulating inflammatory cytokine production by human articular chondrocytes (From the Cell Migration Gateway ITGA5 ITGB1ImmunogenThe original antibody was generated by immunizing BALB/c mice with Nalm-6	Storage	Store at 4C in the dark.
Preservative       0.05% Sodium Azide         Isotype       IgG         Conjugate       Alexa Fluor 405         Purity       Protein A purified         Buffer       50mM Sodium Borate         Product Description       Product Description         Host       Rabbit         Gene ID       3678         Gene Symbol       ITGA5         Species       Human         Specificity/Sensitivity       This antibody binds human integrin alpha 5 beta 1 near the N-terminus. Epitope mapping of SG/19 revealed that Thr82 in the 1 subount, located at the outer face of the boundary between the I-like and hybrid domains, was the key binding determinant for this antibody. Integrin alpha5 beta 1 is also known as fibronectin receptor and binds matrix macromolecules and proteinases and thereby stimulates angiogenesis. Integrin alpha5beta1, is a member of the integrin family, is a heterodimer composed by two subunits, alpha5 (TGA5/CD49e) and beta1 (ITGB1/CD29), and both are necessary for complete biological functions. The interaction of VLA-5 with fibronectin plays an important role in regulating inflammatory cytokine production by human articular chondrocytes (From the Cell Migration Gateway ITGA5 ITGB1         Immunogen       The original antibody was generated by immunizing BALB/c mice with Nalm-6	Clonality	Monoclonal
IsotypeIgGConjugateAlexa Fluor 405PurityProtein A purifiedBuffer50mM Sodium BorateProduct DescriptionHostRabbitGene ID3678Gene SymbolITGA5SpeciesHumanSpecificity/SensitivityThis antibody binds human integrin alpha 5 beta 1 near the N-terminus. Epitope mapping of SG/19 revealed that Thr82 in the 1 subunit, located at the outer face of the boundary between the I-like and hybrid domains, was the key binding determinant for this antibody. Integrin alpha5 beta 1 is also known as fibronectin receptor and binds matrix macromolecules and proteinases and thereby stimulates angiogenesis. Integrin alpha5beta1, is a member of the integrin family, is a heterodimer composed by two subunits, alpha5 (ITGA5/CD49e) and beta1 (ITGB1/CD29), and both are necessary for complete biological functions. The interaction of VLA-5 with fibronectin plays an important role in regulating inflammatory cytokine production by human articular chondrocytes (From the Cell Migration Gateway ITGA5 ITGB1ImmunogenThe original antibody was generated by immunizing BALB/c mice with Nalm-6	Clone	SG/19
ConjugateAlexa Fluor 405PurityProtein A purifiedBuffer50mM Sodium BorateProduct DescriptionHostRabbitGene ID3678Gene SymbolITGA5SpeciesHumanSpecificity/SensitivityThis antibody binds human integrin alpha 5 beta 1 near the N-terminus. Epitope of the boundary between the 1-like and hybrid domains, was the key binding determinant for this antibody. Integrin alpha5 beta 1 is also known as fibronectin receptor and binds matrix macromolecules and proteinases and thereby stimulates angiogenesis. Integrin alpha5beta1, is a member of the integrin family, is a heterodimer composed by two subunits, alpha5 (ITGA5/CD49e) and beta1 (ITGB1/CD29), and both are necessary for complete biological functions. The interaction of VLA-5 with fibronectin plays an important role in regulating inflammatory cytokine production by human articular chondrocytes (From the Cell Migration Gateway ITGA5 ITGB1ImmunogenThe original antibody was generated by immunizing BALB/c mice with Nalm-6	Preservative	0.05% Sodium Azide
PurityProtein A purifiedBuffer50mM Sodium BorateProduct DescriptionHostRabbitGene ID3678Gene SymbolITGA5SpeciesHumanSpecificity/SensitivityThis antibody binds human integrin alpha 5 beta 1 near the N-terminus. Epitope mapping of SG/19 revealed that Thr82 in the 1 subunit, located at the outer face of the boundary between the I-like and hybrid domains, was the key binding determinant for this antibody. Integrin alpha5 beta 1 is also known as fibronectin receptor and binds matrix macromolecules and proteinases and thereby stimulates angiogenesis. Integrin alpha5beta1, is a member of the integrin family, is a heterodimer composed by two subunits, alpha5 (ITGA5/CD49e) and beta1 (ITGB1/CD29), and both are necessary for complete biological functions. The interaction of VLA-5 with fibronectin plays an important role in regulating inflammatory cytokine production by human articular chondrocytes (From the Cell Migration Gateway ITGA5 ITGB1ImmunogenThe original antibody was generated by immunizing BALB/c mice with Nalm-6	Isotype	IgG
Buffer50mM Sodium BorateProduct DescriptionHostRabbitGene ID3678Gene SymbolITGA5SpeciesHumanSpecificity/SensitivityThis antibody binds human integrin alpha 5 beta 1 near the N-terminus. Epitope mapping of SG/19 revealed that Thr82 in the 1 subunit, located at the outer face of the boundary between the I-like and hybrid domains, was the key binding determinant for this antibody. Integrin alpha5 beta 1 is also known as fibronectin receptor and binds matrix macromolecules and proteinases and thereby stimulates angiogenesis. Integrin alpha5beta1, is a member of the integrin family, is a heterodimer composed by two subunits, alpha5 (ITGA5/CD49e) and beta1 (ITGB1/CD29), and both are necessary for complete biological functions. The interaction of VLA-5 with fibronectin plays an important role in regulating inffammatory cytokine production by human articular chondrocytes (From the Cell Migration Gateway ITGA5 ITGB1ImmunogenThe original antibody was generated by immunizing BALB/c mice with Nalm-6	Conjugate	Alexa Fluor 405
Product DescriptionHostRabbitGene ID3678Gene SymbolITGA5SpeciesHumanSpecificity/SensitivityThis antibody binds human integrin alpha 5 beta 1 near the N-terminus. Epitope mapping of SG/19 revealed that Thr82 in the 1 subunit, located at the outer face of the boundary between the I-like and hybrid domains, was the key binding determinant for this antibody. Integrin alpha5 beta 1 is also known as fibronectin receptor and binds matrix macromolecules and proteinases and thereby stimulates angiogenesis. Integrin alpha5beta1, is a member of the integrin family, is a heterodimer composed by two subunits, alpha5 (ITGA5/CD49e) and beta1 (ITGB1/CD29), and both are necessary for complete biological functions. The interaction of VLA-5 with fibronectin plays an important role in regulating inflammatory cytokine production by human articular chondrocytes (From the Cell Migration Gateway ITGA5 ITGB1ImmunogenThe original antibody was generated by immunizing BALB/c mice with Nalm-6	Purity	Protein A purified
HostRabbitGene ID3678Gene SymbolITGA5SpeciesHumanSpecificity/SensitivityThis antibody binds human integrin alpha 5 beta 1 near the N-terminus. Epitope mapping of SG/19 revealed that Thr82 in the 1 subunit, located at the outer face of the boundary between the I-like and hybrid domains, was the key binding determinant for this antibody. Integrin alpha5 beta 1 is also known as fibronectin receptor and binds matrix macromolecules and proteinases and thereby stimulates angiogenesis. Integrin alpha5beta1, is a member of the integrin family, is a heterodimer composed by two subunits, alpha5 (ITGA5/CD49e) and beta1 (ITGB1/CD29), and both are necessary for complete biological functions. The interaction of VLA-5 with fibronectin plays an important role in regulating inflammatory cytokine production by human articular chondrocytes (From the Cell Migration Gateway ITGA5 ITGB1ImmunogenThe original antibody was generated by immunizing BALB/c mice with Nalm-6	Buffer	50mM Sodium Borate
Gene ID3678Gene SymbolITGA5SpeciesHumanSpecificity/SensitivityThis antibody binds human integrin alpha 5 beta 1 near the N-terminus. Epitope mapping of SG/19 revealed that Thr82 in the 1 subunit, located at the outer face of the boundary between the I-like and hybrid domains, was the key binding determinant for this antibody. Integrin alpha5 beta 1 is also known as fibronectin receptor and binds matrix macromolecules and proteinases and thereby stimulates angiogenesis. Integrin alpha5beta1, is a member of the integrin family, is a heterodimer composed by two subunits, alpha5 (ITGA5/CD49e) and beta1 (ITGB1/CD29), and both are necessary for complete biological functions. The interaction of VLA-5 with fibronectin plays an important role in regulating inflammatory cytokine production by human articular chondrocytes (From the Cell Migration Gateway ITGA5 ITGB1ImmunogenThe original antibody was generated by immunizing BALB/c mice with Nalm-6	Product Description	
Gene SymbolITGA5SpeciesHumanSpecificity/SensitivityThis antibody binds human integrin alpha 5 beta 1 near the N-terminus. Epitope mapping of SG/19 revealed that Thr82 in the 1 subunit, located at the outer face of the boundary between the I-like and hybrid domains, was the key binding determinant for this antibody. Integrin alpha5 beta 1 is also known as fibronectin receptor and binds matrix macromolecules and proteinases and thereby stimulates angiogenesis. Integrin alpha5beta1, is a member of the integrin family, is a heterodimer composed by two subunits, alpha5 (ITGA5/CD49e) and beta1 (ITGB1/CD29), and both are necessary for complete biological functions. The interaction of VLA-5 with fibronectin plays an important role in regulating inflammatory cytokine production by human articular chondrocytes (From the Cell Migration Gateway ITGA5 ITGB1ImmunogenThe original antibody was generated by immunizing BALB/c mice with Nalm-6	Host	Rabbit
SpeciesHumanSpecificity/SensitivityThis antibody binds human integrin alpha 5 beta 1 near the N-terminus. Epitope mapping of SG/19 revealed that Thr82 in the 1 subunit, located at the outer face of the boundary between the I-like and hybrid domains, was the key binding determinant for this antibody. Integrin alpha5 beta 1 is also known as fibronectin receptor and binds matrix macromolecules and proteinases and thereby stimulates angiogenesis. Integrin alpha5beta1, is a member of the integrin family, is a heterodimer composed by two subunits, alpha5 (ITGA5/CD49e) and beta1 (ITGB1/CD29), and both are necessary for complete biological functions. The interaction of VLA-5 with fibronectin plays an important role in regulating inflammatory cytokine production by human articular chondrocytes (From the Cell Migration Gateway ITGA5 ITGB1ImmunogenThe original antibody was generated by immunizing BALB/c mice with Nalm-6	Gene ID	3678
Specificity/SensitivityThis antibody binds human integrin alpha 5 beta 1 near the N-terminus. Epitope mapping of SG/19 revealed that Thr82 in the 1 subunit, located at the outer face of the boundary between the I-like and hybrid domains, was the key binding determinant for this antibody. Integrin alpha5 beta 1 is also known as fibronectin receptor and binds matrix macromolecules and proteinases and thereby stimulates angiogenesis. Integrin alpha5beta1, is a member of the integrin family, is a heterodimer composed by two subunits, alpha5 (ITGA5/CD49e) and beta1 (ITGB1/CD29), and both are necessary for complete biological functions. The interaction of VLA-5 with fibronectin plays an important role in regulating inflammatory cytokine production by human articular chondrocytes (From the Cell Migration Gateway ITGA5 ITGB1ImmunogenThe original antibody was generated by immunizing BALB/c mice with Nalm-6	Gene Symbol	ITGA5
<ul> <li>mapping of SG/19 revealed that Thr82 in the 1 subunit, located at the outer face of the boundary between the I-like and hybrid domains, was the key binding determinant for this antibody. Integrin alpha5 beta 1 is also known as fibronectin receptor and binds matrix macromolecules and proteinases and thereby stimulates angiogenesis. Integrin alpha5beta1, is a member of the integrin family, is a heterodimer composed by two subunits, alpha5 (ITGA5/CD49e) and beta1 (ITGB1/CD29), and both are necessary for complete biological functions. The interaction of VLA-5 with fibronectin plays an important role in regulating inflammatory cytokine production by human articular chondrocytes (From the Cell Migration Gateway ITGA5 ITGB1</li> <li>Immunogen</li> </ul>	Species	Human
	Specificity/Sensitivity	mapping of SG/19 revealed that Thr82 in the 1 subunit, located at the outer face of the boundary between the I-like and hybrid domains, was the key binding determinant for this antibody. Integrin alpha5 beta 1 is also known as fibronectin receptor and binds matrix macromolecules and proteinases and thereby stimulates angiogenesis. Integrin alpha5beta1, is a member of the integrin family, is a heterodimer composed by two subunits, alpha5 (ITGA5/CD49e) and beta1 (ITGB1/CD29), and both are necessary for complete biological functions. The interaction of VLA-5 with fibronectin plays an important role in regulating inflammatory cytokine production by human articular chondrocytes (From the
	Immunogen	

www.novusbio.com



	Alexa Fluor (R) products are provided under an intellectual property license from Life Technologies Corporation. The purchase of this product conveys to the buyer the non-transferable right to use the purchased product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components, or any materials made using the product or its components, in any activity to generate revenue, which may include, but is not limited to use of the product or its components: (i) in manufacturing; (ii) to provide a service, information, or data in return for payment; (iii) for therapeutic, diagnostic or prophylactic purposes; or (iv) for resale, regardless of whether they are resold for use in research. For information on purchasing a license to this product for purposes other than as described above, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad, CA 92008 USA or outlicensing@lifetech.com. This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet.
Product Application Details	
Applications	Flow Cytometry, Functional, Immunocytochemistry/ Immunofluorescence, Immunoprecipitation
Recommended Dilutions	Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunoprecipitation, Functional
Application Notes	Optimal dilution of this antibody should be experimentally determined.

Notes





#### Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

#### **Bio-Techne Canada**

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

### **Bio-Techne Ltd**

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

#### **General Contact Information**

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

#### Products Related to NBP3-43579AF405

#### Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP3-43579AF405

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

www.novusbio.com

