# **Product Datasheet**

# MACC1 Antibody (CL0856) - Azide and BSA Free NBP3-43929

Unit Size: 100 ug

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP3-43929

Updated 3/13/2025 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications
Submit a review at www.novusbio.com/reviews/destination/NBP3-43929




## NBP3-43929

MACC1 Antibody (CL0856) - Azide and BSA Free

MACC1 Antibody (CL0856) - Azide and BSA Free	
Product Information	
100 ug	
LYOPH mg/ml	
Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.	
Monoclonal	
CL0856	
No Preservative	
Centrifuge the vial of lyophilized antibody at 12,000 x g for 20 seconds. Reconstitute by adding sterile, distilled water to achieve a final antibody concentration of 1mg/ml.	
IgG1	
Protein A purified	
Lyophilized from a 0.2 um filtered solution in PBS with Trehalose	
Mouse	
346389	
MACC1	
Human	
This antibody was generated using a recombinant protein sequence of Q6ZN28, with the exact immunogen sequence remaining proprietary.	
Immunocytochemistry/ Immunofluorescence, Immunohistochemistry-Paraffin	
Immunocytochemistry/ Immunofluorescence 2-10 ug/ml, Immunohistochemistry- Paraffin 1:500 - 1:1000	
For IHC-Paraffin, HIER pH 6 retrieval is recommended. Immunocytochemistry/Immunofluorescence Fixation Permeabilization: Use PFA/Triton X-100.	


# **Images**

Staining of A-431 cells using the anti-MACC1 monoclonal antibody) .







Analysis in human small intestine and skeletal muscle tissues using NBP3-43929 antibody. Corresponding MACC1 RNA-seq data are presented for the same tissues.




Staining of human kidney shows strong cytoplasmic positivity in cells in tubules.




Staining of human colorectal cancer shows strong cytoplasmic positivity in tumor cells.



Staining of human skeletal muscle shows negative to very weak cytoplasmic positivity in myocytes.



Staining of human small intestine shows strong cytoplasmic positivity in glandular cells.





# Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

#### **Bio-Techne Canada**

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

#### **Bio-Techne Ltd**

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

#### **General Contact Information**

www.novusbio.com
Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

### **Products Related to NBP3-43929**

HAF007 Goat anti-Mouse IgG Secondary Antibody [HRP]

NB720-B Rabbit anti-Mouse IgG (H+L) Secondary Antibody [Biotin]

NBP1-97005-0.5mg Mouse IgG1 Isotype Control (MG1)
NBP1-89352PEP MACC1 Recombinant Protein Antigen

#### Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP3-43929

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

