Product Datasheet

Macro H2A.2 Antibody (CL5263) - Azide and BSA Free NBP3-44246

Unit Size: 100 ug

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP3-44246

Updated 3/16/2025 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP3-44246




NBP3-44246

Macro H2A.2 Antibody (CL5263) - Azide and BSA Free

Macro H2A.2 Antibody (CL5263) - Azide and BSA Free	
Product Information	
100 ug	
LYOPH mg/ml	
Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.	
Monoclonal	
CL5263	
No Preservative	
Centrifuge the vial of lyophilized antibody at 12,000 x g for 20 seconds. Reconstitute by adding sterile, distilled water to achieve a final antibody concentration of 1mg/ml.	
IgG1	
Protein A purified	
Lyophilized from a 0.2 um filtered solution in PBS with Trehalose	
Product Description	
Mouse	
55506	
MACROH2A2	
Human	
This antibody was generated using a synthetic peptide of Q9P0M6, with the exact immunogen sequence remaining proprietary.	
Product Application Details	
Immunocytochemistry/ Immunofluorescence, Immunohistochemistry-Paraffin	
Immunocytochemistry/ Immunofluorescence 2-10 ug/ml, Immunohistochemistry- Paraffin 1:200 - 1:500	
For IHC-Paraffin, HIER pH 6 retrieval is recommended. For Immunocytochemistry/ Immunofluorescence/IF, PFA/Triton X-100 is recommended for fixation/permeabilization.	

Images

Staining of human kidney shows strong nuclear positivity in cells in renal glomeruli and a subset of renal tubules.





Page 2 of 3 v.20.1 Updated 3/16/2025 Staining of human fallopian tube shows strong nuclear immunoreactivity in glandular cells and moderate positivity in stromal cells. Staining of human cerebral cortex shows moderate to strong nuclear immunoreactivity in neuronal and glial cells. Staining of PC-3 cells using the Anti-Macro H2A.2 monoclonal antibody) Staining of human testis shows strong nuclear immunoreactivity in a subset of cells in seminiferous tubules.





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

Products Related to NBP3-44246

HAF007 Goat anti-Mouse IgG Secondary Antibody [HRP]

NB720-B Rabbit anti-Mouse IgG (H+L) Secondary Antibody [Biotin]

NBP1-97005-0.5mg Mouse IgG1 Isotype Control (MG1)

H00055506-P01-10ug Recombinant Human Macro H2A.2 GST (N-Term) Protein

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP3-44246

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

